

On the Mass Eigenstate Purity of ^8B Solar Neutrinos¹

Stephen Parke*, Hiroshi Nunokawa[†] and Renata Zukanovich Funchal**

**Theoretical Physics Department, Fermi National Accelerator Laboratory,
P.O. Box 500, Batavia, IL 60510, USA
email: parke@fnal.gov*

*[†]Departamento de Física, Pontifícia Universidade Católica do Rio de Janeiro,
C. P. 38071, 22452-970, Rio de Janeiro, Brazil
email: nunokawa@fis.puc-rio.br*

***Instituto de Física, Universidade de São Paulo,
C. P. 66.318, 05315-970 São Paulo, Brazil
email: zukanov@if.usp.br*

Abstract. We give a brief report on our recent paper, Ref. [1], in which we calculate the ν_2 mass eigenstate purity of ^8B solar neutrinos as $91 \pm 2\%$.

In a two neutrino analysis, the *day-time* CC/NC of SNO, which is equivalent to the day-time average ν_e survival probability, $\langle P_{ee} \rangle$, is given by

$$\left. \frac{\text{CC}}{\text{NC}} \right|_{\text{day}} = \langle P_{ee} \rangle = f_1 \cos^2 \theta_\odot + f_2 \sin^2 \theta_\odot, \quad (1)$$

where f_1 and $f_2 = 1 - f_1$ are understood to be the ν_1 and ν_2 fractions, respectively, averaged over the ^8B neutrino energy spectrum weighted with the charged current cross section. Therefore the ν_1 fraction (or how much f_2 differs from 100%) is given by

$$f_1 = \frac{\left(\left. \frac{\text{CC}}{\text{NC}} \right|_{\text{day}} - \sin^2 \theta_\odot \right)}{\cos 2\theta_\odot} = \frac{(0.347 - 0.311)}{0.378} \approx 10 \pm ?? \%, \quad (2)$$

where the central values of the recent SNO analysis, [2], have been used. Due to the correlations in the uncertainties between the CC/NC ratio and $\sin^2 \theta_\odot$ we are unable to estimate the uncertainty on f_1 from their analysis. Note, that if the fraction of ν_2 were 100%, then $\frac{\text{CC}}{\text{NC}} = \sin^2 \theta_\odot$.

Using the analytical analysis of the Mikheyev-Smirnov-Wolfenstein (MSW) effect given in Ref. [3], the mass eigenstate fractions are given by

$$f_2 = 1 - f_1 = \langle \sin^2 \theta_\odot^N + P_x \cos 2\theta_\odot^N \rangle_{^8\text{B}}, \quad (3)$$

¹ Presented by Stephen Parke at PANIC05, October 24-28, 2005 in Santa Fe, NM, USA.

^8B : ν_2 fraction (%)




FIGURE 1. The ν_2 fraction (%) in the $\delta m_{\odot}^2$ versus $\sin^2 \theta_{\odot}$ plane. (a) The solid and dashed (blue) lines are the 90, 65, 35 and 10% iso-contours of the fraction of the solar ^8B neutrinos that are ν_2 's. The current best fit value, indicated by the open circle with the cross, is close to the 90% contour. The iso-contour for an electron neutrino survival probability, $\langle P_{ee} \rangle$, equal to 35% is the dot-dashed (red) “triangle” formed by the 65% ν_2 purity contour for small $\sin^2 \theta_{\odot}$ and a vertical line in the pure ν_2 region at $\sin^2 \theta_{\odot} = 0.35$. Except at the top and bottom right hand corners of this triangle the ν_2 purity is either 65% or 100%. (b) Focuses in on the current allowed region. The 68 and 95% CL are shown by the shaded areas with the best fit values indicated by the star using the combined fit of KamLAND and solar neutrino data given in [2].

where $\theta_{\odot}^N$ is the mixing angle defined at the ν_e production point and P_x is the probability of the neutrino to jump from one mass eigenstate to the other during the Mikheyev-Smirnov resonance crossing. The average $\langle \dots \rangle_{^8\text{B}}$ is over the electron density of the ^8B ν_e production region in the center of the Sun predicted by the Standard Solar Model and the energy spectrum of ^8B neutrinos weighted with SNO's charged current cross section. Fig. 1 shows the iso-contours of this averaged ν_2 fraction using a threshold of 5.5 MeV on the kinetic energy of the recoil electrons. Thus the ^8B energy weighted average fraction of ν_2 's observed by SNO is

$$f_2 = 91 \pm 2\% \quad \text{at the 95\% CL.} \quad (4)$$

Hence, the ^8B solar neutrinos are the purest mass eigenstate neutrino beam known so far.

Since the propagation of ^8B neutrinos, in the Sun, is highly adiabatic, the fraction of ν_2 , and consequently, the SNO CC/NC ratio is determined only by the effective value of the matter potential at neutrino production near the center of the Sun. This implies that if we can measure $\sin^2 \theta_{\odot}$ using an experiment independent of the ^8B solar neutrinos, then from the measured value of SNO's CC/NC ratio we can determine the effective value of the matter potential at production. One cannot extract information on the electron number density distribution or the ^8B neutrino production distribution, separately, but only a single characteristic of the convolution of these two distributions, $Y_e \rho|_{eff}^{^8\text{B}}$. Fig. 2 demonstrates this observation and puts a lower bound on $Y_e \rho|_{eff}^{^8\text{B}} > 40 g/cm^3$, see Ref. [1] for details.




FIGURE 2. The iso-contours of $Y_e \rho |_{eff}^{8B}$ in the $\sin^2 \theta_\odot - CC/NC|_{day}$ plane. The line labeled SSM is the Standard Solar Model prediction for $Y_e \rho |_{eff}^{8B}$ ($\approx 85 \text{ g/cm}^3$). SNO's range of observed values of CC/NC are indicated by the shaded horizontal bands.

If $\sin^2 \theta_{13}$ is non-zero then we must use a three neutrino analysis. The three neutrino mass eigenstate fractions, $\mathcal{F}_i$ are

$$\begin{aligned} \mathcal{F}_1 &= f_1 = 0.09 \mp 0.02, \\ \mathcal{F}_2 &= f_2 - \sin^2 \theta_{13} \approx 0.91 \pm 0.02 - \sin^2 \theta_{13}, \\ \mathcal{F}_3 &= \sin^2 \theta_{13}. \end{aligned} \tag{5}$$

A Taylor series expansion about the two flavor mass eigenstate fractions, f_i , has been used since $\sin^2 \theta_{13}$ is known to be less than 0.04.

In summary, the 8B solar neutrinos are produced and propagate from the center of sun to the earth's surface as almost a pure ν_2 mass eigenstate with a purity between 85 and 93%.

REFERENCES

1. H. Nunokawa, S. Parke and R. Z. Funchal, "What fraction of B-8 Solar neutrinos arrive at the Earth as a ν_2 mass eigenstate", Fermilab-Pub-05-049, arXiv:hep-ph/0601198.
2. B. Aharmim *et al.* [SNO Collaboration], arXiv:nucl-ex/0502021.
3. S. J. Parke, Phys. Rev. Lett. **57**, 1275 (1986); S. J. Parke and T. P. Walker, Phys. Rev. Lett. **57**, 2322 (1986).